



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

extent of this region to the length of the elongating zone. PICCARD's method of 1904, which has lately been justified by HABERLANDT,<sup>24</sup> is considered by NEWCOMBE as "too precarious to be satisfactory." All the phenomena, he concludes, "accord equally well with the hypothesis of the extension of the sensitiveness through the elongating zone, but diminishing from the apex backward; or . . . of a more equable sensitiveness through the elongating zone, and a stronger autotropism in the posterior than in the anterior part." We should much prefer the former hypothesis; if for no other reason, because it is unfortunate to postulate "autotropism" when it can be avoided.—C. R. B.

**Development of Marchantia.**—Because no consecutive account of the development of the sexual organs and sporogonium of *Marchantia*, complete in itself, has been published by one author, DURAND, while preparing slides for class use, has published an account of the development illustrated with a close series of figures.<sup>25</sup> The account contributes little which is new to students making a critical study of this form. For the first time, although it has been illustrated by CURTIS, formal attention is called to the familiar "mushroom anchor" foot. One striking feature in the development of the sporophyte has been overlooked: the sterile plate of cells at the apex of the capsule, and also the occasional appearance of a columella, which in some instances extends entirely through the center of the capsule. Because of its relation to the theory of sterilization of sporogenous tissue this plate of cells and the occasional columella should have some attention.—W. J. G. LAND.

**The nucleus of bacteria.**—MEYER claims<sup>26</sup> that the following methods will differentiate a nucleus in the bacteria. The particular form used was *Bacillus Pasteurianus*. First method: boil in water, stain 24 hours in hematoxylin, and differentiate in weak hydrochloric acid. The nuclei of young spores are sharply outlined. Second method: fix in Flemming's solution, harden in 20 per cent. alcohol, stain in Delafield's hematoxylin, and differentiate with hydrochloric acid. Third method: fix in Flemming's solution, harden in alcohol, stain in iron alum hematoxylin, and differentiate under the cover glass with ammonium ferrosulfate. Judging from the figures, this method gives the best results.

In the opinion of the reviewer, the fact that MEYER does not believe that any nucleus has as yet been demonstrated in the Cyanophyceae would not inspire confidence in his interpretation.—CHARLES J. CHAMBERLAIN.

**Anatomy of Sapotaceae.**—Incidentally, in seeking the origin of laticiferous tissue in Sapotaceae, Miss SMITH<sup>27</sup> made anatomical studies of seedlings of fourteen

<sup>24</sup> BOT. GAZETTE 47:482. 1909.

<sup>25</sup> DURAND, ELIAS J., The development of the sexual organs and sporogonium of *Marchantia polymorpha*. Bull. Torr. Bot. Club 35:321-335. pls. 21-25. 1908.

<sup>26</sup> MEYER, ARTHUR, Der Zellkern der Bakterien. Flora 98:335-340. figs. 3. 1908.

<sup>27</sup> SMITH, WINIFRED, The anatomy of some sapotaceous seedlings. Trans. Linn. Soc. London II. 7:189-200. pls. 25, 26. 1909.